



A validated system for ligation-free USER™ -based assembly of expression vectors for mammalian cell engineering

Lund, Anne Mathilde; Kildegaard, Helene Fastrup; Hansen, Bjarne Gram; Holm, Dorte Koefoed; Andersen, Mikael Rørdam; Mortensen, Uffe Hasbro

Publication date:
2013

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Lund, A. M., Kildegaard, H. F., Hansen, B. G., Holm, D. K., Andersen, M. R., & Mortensen, U. H. (2013). *A validated system for ligation-free USER™ -based assembly of expression vectors for mammalian cell engineering*. Poster session presented at 23rd ESACT meeting, Lille, France.
<http://www.lepublicsystemepco.com/events.php?IDManif=640&IDModule=71&IDRub=155>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

A validated system for ligation-free uracil-excision based assembly of expression vectors for mammalian cell engineering


Anne Mathilde Lund^{1,*}, Helene Faustrup Kildegaard², Bjarne Gram Hansen³, Dorte Koefoed Holm¹, Mikael Rørdam Andersen¹, Uffe Hasbro Mortensen¹

(1) Center for Microbial Biotechnology, Department of Systems Biology, Technical University of Denmark, Kgs. Lyngby, Denmark; (2) Current address: Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, Hoersholm, Denmark; (3) Current address: Novozymes A/S, Bagsvaerd, Denmark
(*) amalu@bio.dtu.dk

Introduction

The development in the field of mammalian cell factories require fast and high-throughput methods, this means a high need for simpler and more efficient cloning techniques. For optimization of protein expression by genetic engineering and for allowing metabolic engineering in mammalian cells, a new versatile expression vector system was developed. This vector system applies the ligation-free uracil-excision cloning technique to construct mammalian expression vectors of multiple parts and with maximum flexibility.

Vector assembly platform allows high-throughput generation of genetic constructs



A novel cassette free assembly system was created for solely PCR-based assembly of multiple parts. The simplicity and flexibility of the uracil-excision cloning technique facilitated the adaption of the vector system into high-throughput cloning of a single element or versatile assembly of multiple elements. An expression vector of seven generic elements can be assembled and ready for transfection within three working days.


The assembly method has been tested with >25 generic parts

A library of the most commonly used generic parts for transient and stable plasmid expression in mammalian cells have been created and tested. All parts are ready-to-clone DNA fragments, which allow fast and versatile vector construction with the possibilities of bi-cistronic expression, compartmentalisation, fusion to reporter gene or tags.

Ready-to-clone DNA fragment library


Promoters: <ul style="list-style-type: none"> CMV PGK SV40 	Terminators: <ul style="list-style-type: none"> BGH polyA SV40 polyA hGH polyA 	Targeting signals: <ul style="list-style-type: none"> Nucleus Peroxisomes Mitochondria Plasma membrane ER Medial Golgi Trans Golgi TGN Golgi Secretion signals
Reporters: <ul style="list-style-type: none"> eGFP eYFP eCFP mCherry ccdB SEAP 	Markers: <ul style="list-style-type: none"> Hygromycin Neomycin 	Epitope-tags: <ul style="list-style-type: none"> c-Myc tag HA tag His tag
Other elements: <ul style="list-style-type: none"> IRES ccdB 		

High efficiency one step ligation-free uracil-excision cloning



Uracil-excision is a PCR-based cloning method that employs the USERTM enzyme, a mixture of a Uracil DNA glycosylase and the DNA glycosylase-lyase Endonuclease VIII. The method generates long complementary overhangs that anneal to each other to form a stable hybridisation product that can be used to transform *E. coli* without prior ligation^{1,2}. The method has advantages of permitting additional modification in the assembly process including point mutations, insertion, deletion as well as seamless fusion of DNA fragments³.

Proof-of-concept: Successful protein targeting to organelles



Transient expression was validated in the U-2 OS cell line by a set of vectors for localisation to the major compartments. Expression of fluorescent proteins and protein secretion of secreted alkaline phosphatase⁴ (SEAP) were achieved in HEK293 cells as well as in CHO-K1 cell lines. Furthermore, constructed vectors have successfully been applied for generating stable CHO-K1 cell lines for protein expression by use of selection pressure with Neomycin.

Perspectives

The developed vector assembly system already facilitates high-throughput construction of plasmid DNA for system-wide studies of CHO cell lines and will eliminate bottlenecks within genomic studies of mammalian cells. In the future, the system is planned to include site-specific integration and knockout cassettes.

Acknowledgement

Thanks to the Danish Cancer Society Research Center for providing template plasmids, the U-2 OS and HEK293 cell line. Thanks to Novo Nordisk for kindly providing the CHO-K1 cell line and thanks to the Novo Nordisk Foundation Center for Protein Research for providing the genes of the fluorescent proteins. A special thank to the 2011 iGEM team DTU-Denmark-2. Finally, a thanks to the Core Facilities of Integrated Microscopy at Copenhagen University for use of confocal microscopy.

References

- B.G. Hansen et al., Versatile enzyme expression and characterization system for *Aspergillus nidulans*, with the *Penicillium brevicompactum* polyketide synthase gene from the mycophenolic acid gene cluster as a test case, *Appl Environ Microbiol.* (2011) 77(9):3044-51.
- H.H. Nour-Eldin et al., USER cloning and USER fusion: the ideal cloning techniques for small and big laboratories, *Methods Mol. Biol.* (2010) 643:185-200.
- M. Nørholm, A mutant Pfu DNA polymerase designed for advanced uracil-excision DNA engineering, *BMC Biotechnol.* (2010) 10(1):21.
- S.S. Tate et al., Secreted alkaline phosphatase: an internal standard for expression of injected mRNAs in the *Xenopus* oocyte, *FASEB J.* (1990) 4(2): 227-231.

